

Research Article

Cross-national measurement invariance of the Patient Health Questionnaire-4 (PHQ-4) as a screening measure for depression and anxiety symptoms in 12 Latin American countries

Tomás Caycho-Rodríguez^{1*}, Lindsey W. Vilca², José Ventura-León³, Pablo D. Valencia⁴, Carlos Carbajal-León⁵, Mario Reyes-Bossio⁶, Mariel Delgado-Campusano⁶, Daniel E. Yupanqui-Lorenzo⁷, Claudio Rojas-Jara⁸, Miguel Gallegos^{8,9,10}, Mauricio Cervigni^{10,11}, Pablo Martino¹², Roberto Polanco-Carrasco¹³, Diego Alejandro Palacios¹⁴, Rodrigo Moreta-Herrera¹⁵, Antonio Samaniego-Pinho¹⁶, Marlon Elías Lobos Rivera¹⁷, Andrés Buschiazzi Figares¹⁸, Diana Ximena Puerta-Cortés¹⁹, Ibraín Enrique Corrales-Reyes²⁰, Raymundo Calderón²¹, Olimpia Petzold²², Andrés Camargo²³, Julio Torales^{24,25,26}, Daniela Ferrufino-Borja²⁷, Agueda Muñoz-del-Carpio-Toia²⁸, Jorge Palacios²⁹, Marion K. Schulmeyer²⁷, Cirilo H. Garcia-Cadena³⁰, María José Baeza-Rivera³¹, Camila Salazar-Fernández³², Nicol A. Barria-Asenjo³³, Jesús Ayala-Colqui³⁴

[1] Faculty of Psychology, Universidad Científica del Sur, Lima, Peru.

[2] Universidad Señor de Sipán, Chiclayo, Perú.

[3] Faculty of Health Sciences, Universidad Privada del Norte, Lima, Perú.

[4] Coordination of Open University and Digital Education, National Autonomous University of Mexico, Mexico City, Mexico.

[5] School of Psychology, Faculty of Communication Sciences, Tourism and Psychology, Universidad de San Martín de Porres, Lima, Perú.

[6] Faculty of Psychology, Peruvian University of Applied Sciences, Lima, Peru.

[7] School of Psychology, University of Sciences and Humanities, Lima, Peru.

- [8] Faculty of Health Sciences. Department of Psychology, Catholic University of Maule, Talca, Chile.
- [9] Postgraduate Program in Psychology. Catholic University of Minas Gerais, Belo Horizonte, Brazil.
- [10] National Scientific and Technical Research Council, Buenos Aires, Argentina.
- [11] Faculty of Psychology, National University of Rosario, Rosario, Argentina.
- [12] Laboratory of Research in Behavioral Sciences (LICIC), Faculty of Psychology, National University of San Luis, San Luis, Argentina.
- [13] Neuropsychology Notebooks, Rancagua, Chile.
- [14] Human Development Center, Mariano Gálvez University, Guatemala, Guatemala
- [15] School of Psychology, Pontifical Catholic University of Ecuador, Ambato, Ecuador.
- [16] Psychology Degree, Faculty of Philosophy, National University of Asuncion, Asuncion, Paraguay.
- [17] School of Psychology, Faculty of Social Sciences, Technological University of El Salvador, San Salvador, El Salvador.
- [18] Alfred Adler Institute Uruguay, Center for Adlerian Studies, Montevideo, Uruguay. [19] Psychology Program, University of Ibagué, Ibagué, Colombia.
- [20] Maxillofacial Surgery Service, Carlos Manuel de Céspedes General University Hospital, Granma University of Medical Sciences, Bayamo, Granma, Cuba,
- [21] State College of Psychologists in Intervention of Jalisco A.C. Guadalajara, Jalisco, Mexico.
- [22] Lone Star College, Conroe, The United States of America.
- [23] School of Health and Sport Sciences, Fundación Universitaria del Área Andina, Bogotá, Colombia.
- [24] Chair of Medical Psychology, Faculty of Medical Sciences, Universidad Nacional de Asunción, San Lorenzo, Paraguay.
- [25] Regional Institute for Health Research, Universidad Nacional de Caaguazú, Coronel Oviedo, Paraguay.
- [26] Faculty of Medical Sciences, Universidad Sudamericana, Salto del Guairá, Paraguay.
- [27] Center for Psychological Research and Counseling, Faculty of Humanities, Communication and Arts, Private University of Santa Cruz de la Sierra, Santa Cruz, Bolivia.
- [28] Universidad Católica de Santa María, Arequipa, Perú.
- [29] Psychology Degree, Faculty of Health Sciences, University of the Valley of Mexico, Mexico City, Mexico.

- [30] School of Psychology, Autonomous University of Nuevo Leon, Monterrey, Mexico.
- [31] Laboratory of Interactions, Culture and Health, Department of Psychology, Faculty of Health Sciences, Universidad Católica de Temuco, Chile Chile.
- [32] Data Analysis Department, Universidad Autónoma de Chile, Santiago, Chile.
- [33] Department of Social Sciences, University of Los Lagos, Osorno, Chile.
- [34] Technological University of Peru, Lima, Peru.

Abstract

The Patient Health Questionnaire-4 (PHQ-4) is a widely used instrument to measure symptoms of depression and anxiety in general, but not much is known about its cross-cultural utility in Latin America. The present study evaluated the measurement invariance of the PHQ-4 in adult samples ($N = 5441$) from 12 Latin American countries (Paraguay, El Salvador, Chile, Ecuador, Colombia, Uruguay, Peru, Mexico, Cuba, Guatemala, Argentina, and Bolivia). The two-factor structure presented a superior fit to the one-factor structure. This provides evidence to support a two-dimensional model of anxious and depressive symptomatology of the PHQ-4, with a good estimate of reliability. The configural, metric, scalar and strict invariance of the PHQ-4 was supported, suggesting that participants from all countries attribute the same meaning to the latent constructs anxiety and depression and their scores can be compared. Additionally, support was provided for the hypothesis that the presence of depressive symptoms significantly predicts subjective well-being. However, the presence of anxiety symptoms fails to significantly predict participants well-being. It is concluded that the PHQ-4 provides a largely comparable measure of anxiety and depression symptoms in a large sample of Latin American countries.

Keywords: Anxiety; depression; invariance; Latin America; Patient Health Questionnaire-4.

Table of Contents

Method
Results
Discussion
References

Psychological Thought, 2024, Vol. 17(2), 450-482, <https://doi.org/10.37708/psyct.v17i2.931>

Received: 2023-12-21. Accepted: 2024-10-01. Published (VoR): 2024-10-31.

Handling Editor: Natasha Angelova, South-West University "Neofit Rilski", Blagoevgrad, Bulgaria. *Corresponding author at: Facultad de Psicología, Universidad Científica del Sur, Lima, Peru.

E-mail: tppcaycho@gmail.com



This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Mental health is an essential aspect of global health. Its importance worldwide has grown, as mental health disorders have become the leading cause of disability, accounting for one of every six years lived with disability ([World Health Organization, 2022](#)). In Latin America, these disorders have experienced an increase in the last decade, which has been followed by a wide gap in their treatment. This situation is critical, as mental health disorders increase the risk of developing other diseases and contribute to intentional and unintentional injuries ([Pan American Health Organization, 2023](#)).

The growth in the prevalence of mental health disorders has been intensified by the COVID-19 pandemic. Latin America, in particular, has shown regional variations in mental health ([Zhang et al., 2022](#)). Before the pandemic of COVID-19, the prevalence rate of severe mental disorders in the Americas region ranged from 2% to 10% ([Kohn et al., 2018](#)). Specifically, annual prevalence rates of anxiety disorders and depression were reported to be 5.8% and 4.9% in the adult population ([Caldas de Almeida & Horvitz-Lennon, 2010](#)). A systematic review and meta-analysis study indicated that, during the pandemic, the prevalences of anxiety and depression increased to 35% and 35%, respectively, with South American countries having a higher prevalence of mental health symptoms compared to Central America (36% versus 28%, $p < 0.001$) ([Zhang et al., 2022](#)). However, symptoms of anxiety (LOR = -0.33; 95 % CI: -0.54, -0.12) and depression (LOR = -0.12; 95 % CI: -0.21, -0.04) were reported to have decreased across the pandemic worldwide ([Cenat et al., 2022](#)). In Latin America, it has been suggested that mental health during the pandemic had a complex dynamic, where the majority were resilient, but there were also vulnerable groups (younger, female and with pre-existing psychiatric diagnoses) who were more likely to deteriorate in mental health over time ([Fernández et al., 2022](#)).

Depressive and anxiety disorders are considered to be two of the leading causes of disability among people aged 15 to 49 years ([Errazuriz & Crisostomo, 2021](#)). However, in primary health care, detection rates for depression and anxiety are less than 50%, which generates a large number of undiagnosed and untreated patients ([Christodoulaki et al., 2022](#)). Appropriate screening for symptoms of anxiety and depression has been recommended as a step prior to initiating treatment, which is associated with improved patient outcomes ([Clark, 2011](#); [Katon & Roy-Byrne, 2007](#)). Different instruments have been developed to improve the detection rates of depression and anxiety symptoms, such as the Beck Depression Inventory (BDI; [Beck et al., 1961](#)), the Hospital Anxiety and Depression Scale (HADS; [Zigmond & Snaith, 1983](#)) or the Depression Anxiety Stress Scale (DASS; [Parkitny & McAuley, 2010](#)). Even though these have

shown evidence of reliability and validity in the Latin American context (Hinz et al., 2014; Lopez et al., 2021; Mautong et al., 2021; Ruiz et al., 2017), their length generates greater time consumption and difficulties for their use by non-specialists (Christodoulaki et al., 2022). This has led to little use of these measures in clinical practice in primary care (Gilbody et al., 2002). In view of this, for some years now there have been brief measures aimed at improving the reliability and efficiency of the detection of anxiety and depression symptoms (Khubchandani et al., 2016).

An ultra-low screening measure is the Patient Health Questionnaire-4 (PHQ-4; Kroenke et al., 2009) composed of four items, which has demonstrated high sensitivity and specificity for detecting depression and anxiety (Stanhope, 2016). Of the four items, two measure the frequency of depressive symptoms (depressed mood and loss of interest), and the other two measure the frequency of anxiety symptoms (nervousness and uncontrollable worry). The PHQ-4 has been validated in different languages and countries worldwide (Christodoulaki et al., 2022; Ghaheri et al., 2020; Kazlauskas et al., 2023; Kim et al., 2021; Kliem et al., 2016; Kocalevent et al., 2014; López Guerra et al., 2022; Löwe et al., 2010; Materu et al., 2020; Wicke et al., 2022). Psychometric studies with the PHQ-4, conducted in different countries and sociodemographic groups, generally supported the presence of a two-factor structure (anxiety and depression), an adequate estimate of reliability and evidence of measurement invariance (MI) across different age groups, genders or others (Ahmadi et al., 2019; Christodoulaki et al., 2022; Ghaheri et al., 2020; Kazlauskas et al., 2023; Khubchandani et al., 2016; Kim et al., 2021; Kocalevent et al., 2014; Larionow & Mudło-Głagolska, 2023; Lenz & Li, 2022; López Guerra et al., 2022; Löwe et al., 2010; Mendoza et al., 2022; Mills et al., 2015; Renovanz et al., 2019). Although the presence of two factors was constant, it has also been suggested that this factor structure would not be completely adequate (Kim et al., 2021). A study in Tanzania suggested that the four items of the PHQ-4 were grouped into a single factor (Materu et al., 2020).

Likewise, there is evidence of the relationship of PHQ-4 scores with lower levels of positive affects (Mendoza et al., 2022), meaning in life (Fong et al., 2023), meaning in life (López Guerra et al., 2022), well-being (Ghaheri et al., 2020) and quality of life (Renovanz et al., 2019), among others. In addition, the PHQ-4 was related to high levels of stress and negative affects (Mendoza et al., 2022), increased frequency of suicidal ideation (Fong et al., 2023), poor sleep quality, subjective fatigue and lack of concentration (Lahlouh & Mustafa, 2020), psychological inflexibility (López Guerra et al., 2022), hopelessness and distress (Kocalevent et al., 2014; Mills et al., 2015).

Even though to date only two studies have validated the PHQ-4 for samples from Colombia (Kocalevent et al., 2014) and Ecuador (López Guerra et al., 2022), their findings cannot be generalized to the entire Latin American context, given the cultural differences. It has been suggested that culture and social contexts determine how people conceptualize and express emotional problems (Ali et al., 2022). The use of the PHQ-4 in samples from different countries, from Latin America and other continents, has further revealed the importance of this questionnaire, but has also raised concerns about measurement invariance (MI). There is a limited number of studies on the MI of the PHQ-4 across countries. Many researchers assume that when they administer the PHQ-4 in different countries, they are assessing the same construct in all samples in the same way and assume that the questionnaire is invariant across groups (Byrne, 2016). However, differences in prevalence rates of anxiety and depression between countries do not necessarily represent real differences. Even if the same symptoms appear in all countries, methodological problems may lead to false conclusions about cross-cultural differences (Bowden & FoxRushby, 2003; van de Vijver & Tanzer, 2004). Cross-country comparisons require instruments with guaranteed comparability of measurements before reliable conclusions can be drawn (Boer et al., 2018). To our knowledge, only one previous study evaluated the MI of the PHQ-4 in seven European countries (Austria, Croatia, Georgia, Germany, Lithuania, Portugal and Sweden) which indicated the presence of partial scalar measurement invariance across countries (Kazlauskas et al., 2023).

MI refers to the equivalence of measurement in different conditions, such as different countries (Baumgartner & Steenkamp, 1998), and is considered as a property of a measuring instrument (Davidov et al., 2014). Therefore, MI makes it possible to examine whether instruments can measure the same construct in different conditions or observed groups (Horn & McArdle, 1992). This implies that the same construct definition and behaviors are presented in the same way in each country. In this sense, it is necessary to have instruments that are cross-culturally invariant in samples from different countries, and that are short and easy to apply, to evaluate symptoms of depression and anxiety. Therefore, the present study aimed to examine the evidence of validity based on the internal structure, based on the relationship with other variables, reliability and cross-country MI of the PHQ-4 in samples from 12 Latin American countries. Based on the only study of cross-national invariance of the PHQ-4 (Kazlauskas et al., 2023), it was hypothesized that the PHQ-4 has a two-factor structure, which provides reliable scores and is invariant across countries. Also, due to evidence relating both depressive, anxious symptoms and well-being (Kinderman et al., 2015; Malone & Wachholtz, 2018, Topp et al., 2015), it was hypothesized that

anxiety and depressive symptoms, as measured by the PHQ-4, negatively predict well-being. This would provide evidence of validity based on the relationship with other variables.

This multinational study would provide information on the comparability of PHQ-4 data obtained in different countries. Giving evidence of the MI of the PHQ-4 is important for an objective comparison of means between groups from different countries. The lack of evidence of MI would suggest that the differences in anxiety and depression means between countries reflect differences in measurement errors and not the true differences between groups. In other words, only when the PHQ-4 items are invariant will we be certain that the differences in anxiety and depression scores in the Latin American context are due to differences between countries and not to measurement errors. Furthermore, the inclusion of several Latin American countries will not only support the universality of the PHQ-4 but will also allow for more rigorous cross-cultural studies.

Method

Participants

A total of 5441 people from 12 Latin American countries (Paraguay, El Salvador, Chile, Ecuador, Colombia, Uruguay, Peru, Mexico, Cuba, Guatemala, Argentina and Bolivia) were selected through a non-probabilistic convenience sample according to the following inclusion criteria: 1) being of legal age, 2) being a resident in each of the participating countries, 3) providing informed consent to participate in the study. The number of participants was determined by using the [Soper \(2023\)](#). The number of observed ($n = 4$) and latent ($n = 2$) variables in the model, the anticipated effect size ($\lambda = 0.3$), the desired probability ($\alpha = 0.05$) and the statistical power level ($1 - \beta = 0.95$) were indicated. Based on these data, a recommended number of participants of 400 individuals was suggested. Although there are countries with a sample size smaller than recommended, another study has suggested that sample sizes in the range of 100 to 200 may be sufficient to adequately estimate the model parameters. Therefore, the sample sizes of all countries are adequate ([Meade & Lautenschlager, 2004](#)).

Table 1 indicates the highest mean age belongs to participants from Guatemala ($M = 40.9$; $SD = 11.9$ years) and Argentina ($M = 42.3$; $SD = 15.2$ years); while, the lowest mean age belongs to participants from Cuba ($M = 24.6$; $SD = 6.8$ years) and Ecuador ($M = 29$; $SD = 10.6$ years). The majority of participants in all countries are women. In addition, there is a higher proportion of married people in most countries ($> 50\%$). Regarding educational level, the majority of participants completed university studies, except in El Salvador (29.6%), Colombia (37.7%), Uruguay (45%),

Cuba (42.7%) and Argentina (49.7%), where the proportion is much lower than 50%. On the other hand, although most of the participants have a permanent job, there is a significant group of them in all countries who are unemployed (16.9% - 51.2%). It can also be seen that the majority of participants reside in an urban area (> 70%). Further details of the sociodemographic characteristics of the participants in each country are shown in Table 1.

Instruments

Sociodemographic data. A form was prepared specifically for this study to obtain information on age, sex, marital status, educational level, area of residence and type of work in each participating country.

The Patient Health Questionnaire-4 (PHQ-4; Kroenke et al., 2009). The PHQ-4 detects the frequency of the main symptoms of a depression and anxiety disorder during the last 2 weeks. It consists of four items, combining the two items of the PHQ-2 (Löwe et al., 2005) and two items of Generalized Anxiety Disorder Scale- 2 items (GAD-2; Kroenke et al., 2007). The first two items measure the frequency of depressive symptoms ("Little interest or pleasure in doing things" and "Feeling depressed or hopeless") and the last two items measure the frequency of anxiety symptoms ("Feeling nervous, anxious or on edge" and "Not being able to stop or control worry"). Each of the PHQ-4 items has four Likert-type response options, ranging from 0 = not at all to 3 = almost every day. The total score ranges from 0 to 12, where higher scores indicate a higher frequency of depressive and anxiety symptoms. For this study, the Spanish version was used, validated in Ecuador by López Guerra et al. (2022). Reliability was optimal for the total scale ($\alpha = 0.879$; $\omega = 0.880$) and its two factors: anxiety ($\alpha = 0.838$; $\omega = 0.838$) and depression ($\alpha = 0.779$; $\omega = 0.780$).

WHO-5 well-being index (WHO-5; World Health Organization, 1998). The WHO-5 evaluates general subjective well-being during the last 2 weeks. It consists of 5 items: (1) "I have felt cheerful and in good spirits", (2) "I have felt calm and relaxed", (3) "I have felt active and energetic", (4) "I have woken up feeling good and rested"]; (5) "My daily life has had interesting things for me". Each item has four Likert-type response options, ranging from "0 = never" to "3 = always". The sum of the scores for each item generates a total WHO-5 score ranging from 0 to 15, where higher scores would indicate greater subjective well-being. For this study, we used the WHO-5 version, which has demonstrated evidence of validity and MI in general population samples from a number of Latin American countries which showed adequate reliability indices in each of the countries evaluated ($\alpha \geq 0.94$; $\omega \geq 0.77$) (Caycho-Rodríguez et al., 2023)

Table 1.
Sociodemographic characteristics of the participants

Sociodemographic Data	Paraguay (<i>n</i> = 918)	El Salvador (<i>n</i> = 741)	Chile (<i>n</i> = 555)	Ecuador (<i>n</i> = 477)	Colombia (<i>n</i> = 414)	Uruguay (<i>n</i> = 322)
Age (M ± SD)	31.4 ± 10.8	29.2 ± 8.8	36.4 ± 11.8	29 ± 10.6	30 ± 12.8	38.3 ± 13.7
Sex, <i>n</i> (%)						
Male	219 (23.9%)	279 (37.7%)	135 (24.3%)	144 (30.2%)	111 (26.8%)	85 (20.9%)
Female	699 (76.1%)	462 (62.3%)	420 (75.7%)	333 (69.8%)	303 (73.2%)	322 (79.1%)
Marital Status <i>n</i> (%)						
Married	603 (65.7%)	543 (73.3%)	266 (47.9%)	328 (68.8%)	298 (72%)	188 (46.2%)
Cohabitant	209 (22.8%)	130 (17.5%)	155 (27.9%)	97 (20.3%)	66 (15.9%)	89 (21.9%)
Divorced	27 (2.9%)	12 (1.6%)	43 (7.7%)	34 (7.1%)	15 (3.6%)	44 (10.8%)
Single	72 (7.8%)	54 (7.3%)	87 (15.7%)	14 (2.9%)	29 (7%)	80 (19.7%)
Widowed	7 (.8%)	2 (.3%)	4 (.7%)	4 (.8%)	6 (1.4%)	6 (1.5%)
Level of Education, <i>n</i> (%)						
Complete university	488 (53.2%)	219 (29.6%)	370 (66.7%)	245 (51.4%)	156 (37.7%)	183 (45%)
Incomplete university	308 (33.6%)	296 (39.9%)	107 (19.3%)	135 (28.3%)	129 (31.2%)	120 (29.5%)
Technical studies complete	20 (2.2%)	34 (4.6%)	44 (7.9%)	12 (2.5%)	39 (9.4%)	42 (10.3%)
Technical studies incomplete	1 (.1%)	9 (1.2%)	8 (1.4%)	3 (.6%)	7 (1.7%)	1 (.2%)
High school complete	80 (8.7%)	111 (15%)	22 (4%)	75 (15.7%)	72 (17.4%)	35 (8.6%)
Secondary incomplete	17 (1.9%)	49 (6.6%)	3 (.5%)	6 (1.3%)	6 (2.4%)	25 (6.1%)
Primary school complete	4 (.4%)	12 (1.6%)	1 (.2%)	1 (.2%)	1 (.2%)	1 (.2%)
Primary incomplete	0 (0%)	11 (1.5%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Area de residence, <i>n</i> (%)						
Urban	811 (88.3%)	584 (78.8%)	482 (86.8%)	358 (75.1%)	383 (92.5%)	393 (96%)
Rural	107 (11.7%)	157 (21.2%)	73 (13.2%)	119 (24.9%)	31 (7.5%)	14 (3.4%)
Type of work, <i>n</i> (%)						
Permanent job	509 (55.4%)	395 (53.3%)	319 (57.5%)	176 (36.9%)	134 (32.4%)	289 (71%)
Temporary job	156 (17%)	92 (12.4%)	81 (14.6%)	72 (15.1%)	68 (16.4%)	26 (6.4%)
Unemployed	253 (27.6%)	254 (32.4%)	155 (27.9%)	229 (48%)	212 (51.2%)	92 (22.6%)
Sociodemographic Data	Perú (<i>n</i> = 371)	México (<i>n</i> = 343)	Cuba (<i>n</i> = 378)	Guatemala (<i>n</i> = 342)	Argentina (<i>n</i> = 316)	Bolivia (<i>n</i> = 264)
Age (M ± SD)	31.7 ± 10.9	33.1 ± 13.3	24.6 ± 6.8	40.9 ± 11.9	42.3 ± 15.2	39.1 ± 14.3
Sex, <i>n</i> (%)						
Male	119 (32.1%)	116 (33.8%)	127 (37.1%)	121 (35.2%)	68 (21.5%)	76 (28.8%)
Female	252 (67.9%)	227 (66.2%)	215 (62.9%)	223 (64.8%)	248 (78.5%)	188 (71.2%)
Marital Status <i>n</i> (%)						
Married	241 (65%)	191 (55.7%)	226 (66.1%)	139 (40.4%)	148 (46.8%)	136 (51.5%)



Cohabitant	70 (18.9%)	113 (32.9%)	46 (13.5%)	151 (43.9%)	89 (28.2%)	82 (31.1%)
Divorced	17 (4.6%)	19 (5.5%)	14 (4.1%)	28 (8.1%)	29 (9.2%)	32 (12.1%)
Single	41 (11.1%)	15 (4.4%)	55 (16.1%)	20 (5.8%)	41 (13%)	9 (3.4%)
Widowed	2 (.5%)	5 (1.5%)	1 (.3)	6 (1.7%)	9 (2.8%)	5 (1.9%)
Level of Education, <i>n</i> (%)						
Complete university	216 (58.2%)	178 (51.9%)	146 (42.7%)	208 (60.5%)	157 (49.7%)	173 (65.5%)
Incomplete university	105 (28.3%)	95 (27.7%)	172 (50.3%)	74 (21.5%)	87 (27.5%)	63 (23.9%)
Technical studies complete	21 (5.7%)	45 (13.1%)	11 (3.25)	22 (6.4%)	32 (10.1%)	15 (5.7%)
Technical studies incomplete	8 (2.2%)	4 (1.2%)	2 (.6%)	6 (1.7%)	4 (1.3%)	0 (0%)
High school complete	16 (4.3%)	20 (5.8%)	5 (1.5%)	23 (6.7%)	28 (8.9%)	7 (2.7%)
Secondary incomplete	4 (1.1%)	1 (.3%)	6 (1.8%)	10 (2.9%)	4 (1.3%)	5 (1.9%)
Primary school complete	0 (0%)	0 (0%)	0 (0%)	1 (.3%)	4 (1.3%)	1 (.4%)
Primary incomplete	1 (.3%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Area de residence, <i>n</i> (%)						
Urban	329 (88.7%)	320 (93.3%)	301 (88%)	320 (93%)	300 (94.9%)	255 (96.6%)
Rural	42 (11.3%)	23 (6.7%)	41 (12%)	24 (7%)	16 (5.1%)	9 (3.4%)
Type of work, <i>n</i> (%)						
Permanent job	155 (41.8%)	162 (47.2%)	237 (69.3%)	232 (67.4%)	203 (64.2%)	110 (41.7%)
Temporary job	77 (20.8%)	64 (18.7%)	15 (4.4%)	54 (15.7%)	43 (13.6%)	61 (23.1%)
Unemployed	139 (37.5%)	117 (34.1%)	90 (26.3%)	58 (16.9%)	70 (22.2%)	93 (35.2%)



Procedure

An online questionnaire was developed through the Google Forms®. The questionnaire was divided into three parts. In the first part, the objectives of the study and the informed consent were presented. Only participants who gave informed consent agreed to the remaining two parts. In the second part, the sociodemographic card was presented, and finally, in the third part, the PHQ-4 and WHO-5 were presented. The online questionnaire was distributed through social networks, such as Facebook, Instagram, WhatsApp, and email. The data collection procedure was the same in each of the 12 participating countries.

All recommendations for reporting results of questionnaires and online surveys were followed (CHERRIES) (Eysenbach, 2004; López-Rodríguez, 2019). The study protocol was reviewed and approved by the Department of Medical Psychology of the National University of Asuncion (Paraguay; approval number: 001_011_2023), which ensured compliance with the ethical aspects of this type of research, such as anonymity and voluntary participation, confidentiality of the data and the possibility of leaving the study whenever they wished.

Data analysis

The factor structure of the PHQ-4 scale was evaluated through a Confirmatory Factor Analysis (CFA), for which the Diagonally Weighted Least Squares with Mean and Variance corrected (WLSMV) estimator was used since the items presented less than five response categories (Li, 2016). It is essential to mention that a polychoric correlation matrix was calculated to perform the CFA since the items were ordinal. The following indexes were used to evaluate the fit of the models studied: Root mean squared error of approximation (RMSEA), Standardized root mean square residual (SRMR), Comparative fit index (CFI) and Tucker Lewis index (TLI). For the RMSEA and SRMR indexes values less than .08 were considered acceptable (Kline, 2016). For the CFI and TLI indexes, values greater than .95 were considered adequate (Schumacker & Lomax, 2015). Regarding the reliability of the scale, the omega coefficient (McDonald, 1999) and Alpha Cronbach (Cronbach, 1951) was used, where a value greater than .70 is considered adequate (Viladrich et al., 2017). The method developed by Feldt et al. (1987) was used to calculate possible differences in the level of reliability.

To evaluate the factorial invariance of the scale according to the country to which the participants belong, Multi-group Confirmatory Factor Analysis (MGCFAs) was used, where a sequence of hierarchical variance models was proposed: (1) configural invariance, (2) metric invariance, (3) scalar invariance and (4) strict invariance. To compare the sequence of models, the differences



in the CFI (Δ CFI) were used, where values less than $<.010$ show the invariance of the model between groups (Chen, 2007). The RMSEA (Δ RMSEA) was also used, where differences less than $<.015$ evidenced the invariance of the model between groups (Chen, 2007).

Regarding the validity of the scale in relation to other variables, an SEM model was proposed. In this model, the PHQ-4 scale has a significant impact on the level of satisfaction with life. The WLSMV estimator was used to estimate the model and the same adjustment indicators used in the CFA were taken into account.

All statistical analyses were performed using the "lavaan" package (Rosseel, 2012) for the AFC and the SEM model. The package "semTools" (Jorgensen et al., 2018) was used to estimate factorial invariance. Also, the "cocron" library was used to calculate the differences in reliability between the countries (Diedenhofen & Musch, 2016). In all cases, the RStudio environment (RStudio Team, 2018) for R (R Core Team, 2019) was used.

Results

Descriptive Analysis

Table 2 shows that the participants answered all the response categories of the four items. However, the majority of respondents in all countries chose the first two response categories and few chose the higher response categories. It is also observed that in the polychoric correlation matrix all items correlate positively and significantly with each other ($p < .01$), this result occurs in all countries. These results are important, since they suggest the presence of groups of intercorrelated items.

Table 2.*Descriptive analysis of the items in the studied countries*

Region - Country	Ítems	Pattern of response				Polycorrelation matrix			
		C1	C2	C3	C4	PHQ1	PHQ2	PHQ3	PHQ4
Paraguay (n = 918)	PHQ1	40.3%	34.6%	15.1%	9.9%	1			
	PHQ2	45.3%	30.9%	14.5%	9.3%	.84	1		
	PHQ3	47.5%	27.2%	14.6%	10.7%	.74	.74	1	
	PHQ4	49%	27.9%	13.8%	9.3%	.69	.70	.88	1
El Salvador (n = 741)	PHQ1	39.3%	31%	17.8%	11.9%	1			
	PHQ2	39.5%	35.4%	13.8%	11.3%	.84	1		
	PHQ3	42.9%	26.5%	16.7%	13.9%	.74	.75	1	
	PHQ4	42.6%	31.6%	14.4%	11.3%	.70	.73	.86	1
Chile (n = 555)	PHQ1	20.4%	36.9%	24.3%	18.4%	1			
	PHQ2	31.5%	34.4%	19.5%	14.6%	.87	1		
	PHQ3	32.4%	34.2%	18.2%	15.1%	.70	.70	1	
	PHQ4	28.5%	35.7%	19.5%	16.4%	.69	.64	.86	1
Ecuador (n = 477)	PHQ1	35%	35.4%	19.7%	9.9%	1			
	PHQ2	37.5%	36.1%	17.8%	8.6%	.86	1		
	PHQ3	32.3%	36.7%	19.9%	11.1%	.74	.76	1	
	PHQ4	36.5%	32.1%	20.8%	10.7%	.70	.73	.88	1
Colombia (n = 414)	PHQ1	37.2%	33.1%	19.8%	9.9%	1			
	PHQ2	41.5%	33.8%	13.5%	11.1%	.87	1		
	PHQ3	42.3%	27.3%	20%	10.4%	.68	.71	1	
	PHQ4	46.4%	27.5%	15.2%	10.9%	.67	.71	.88	1
Uruguay (n = 407)	PHQ1	37.6%	40%	14.3%	8.1%	1			
	PHQ2	47.4%	36.9%	10.1%	5.7%	.81	1		
	PHQ3	53.1%	29.5%	11.1%	6.4%	.68	.72	1	
	PHQ4	52.3%	29.7%	11.5%	6.4%	.60	.62	.82	1
Perú (n = 371)	PHQ1	34.2%	38%	19.9%	7.8%	1			
	PHQ2	42.6%	32.1%	15.9%	9.4%	.84	1		
	PHQ3	39.6%	34.5%	18.1%	7.8%	.72	.77	1	



Region - Country	Items	Pattern of response				Polycorrelation matrix			
		C1	C2	C3	C4	PHQ1	PHQ2	PHQ3	PHQ4
México (n = 343)	PHQ4	42%	32.9%	16.2%	8.9%	.66	.73	.84	1
	PHQ1	27.1%	39.9%	19.8%	13.1%	1			
	PHQ2	35.9%	35%	16.9%	12.2%	.85	1		
	PHQ3	34.4%	29.7%	22.4%	13.4%	.71	.70	1	
Cuba (n = 342)	PHQ4	37.3%	31.5%	17.2%	14%	.66	.65	.84	1
	PHQ1	46.8%	31.3%	9.6%	12.3%	1			
	PHQ2	47.4%	31.9%	10.2%	10.5%	.83	1		
	PHQ3	52%	27.8%	11.4%	8.8%	.65	.67	1	
Guatemala (n = 344)	PHQ4	49.1%	29.8%	12.6%	8.5%	.58	.57	.76	1
	PHQ1	35.8%	34.3%	18%	11.9%	1			
	PHQ2	45.1%	27%	19.2%	8.7%	.86	1		
	PHQ3	48%	28.8%	13.7%	9.6%	.75	.78	1	
Argentina (n = 316)	PHQ4	49.1%	30.2%	13.4%	7.3%	.72	.77	.88	1
	PHQ1	27.5%	45.9%	14.9%	11.7%	1			
	PHQ2	38.3%	38.6%	12.7%	10.4%	.86	1		
	PHQ3	38.3%	37.3%	13.3%	11.1%	.73	.70	1	
Bolivia (n = 264)	PHQ4	39.2%	37.3%	12.7%	10.8%	.69	.64	.86	1
	PHQ1	23.9%	40.5%	20.8%	14.8%	1			
	PHQ2	28.4%	43.6%	18.2%	9.8%	.85	1		
	PHQ3	29.2%	41.3%	15.2%	14.4%	.70	.79	1	
	PHQ4	34.1%	36%	16.7%	13.3%	.61	.69	.80	1

Note: C1 = None; C2 = Several days; C3 = More than half of the days; C4 = Almost every day.



Validity based on the internal structure and reliability of the scale

Table 3 shows that the original model with two related factors (model 1) demonstrates adequate adjustment indexes in all countries. In addition, it is observed that all the items present high factorial weights in the factor to which they belong. However, the relationship between both factors was found to be high (.77 - .87). In view of this, the performance of a unidimensional model (model 2) for the PHQ-4 scale was proposed. This model showed poor fit indices to the data in all countries. Therefore, model 1 proved to be the model that best explains the factorial structure of the scale and will therefore be used in the following psychometric analyses.

Table 3.
Fit indices of the different scale models in the countries of the Americas

Country	Adjustment index							Factorial weight (λ)				Φ	Reliability			
	χ^2	df	p	CFI	TLI	SRMR	RMSEA [90%CI]	An1	An2	De3	De4	F1~F2	F1 (α/ω)	F2 (α/ω)		
Model 1																
Paraguay ($n = 918$)	.004	1	.950	1.00	1.00	.005	.000 [.000 – .003]	.91	.92	.97	.91	.84	.86	.86	.89	.89
El Salvador ($n = 741$)	.639	1	.424	1.00	1.00	.076	.000 [.000 – .090]	.90	.93	.95	.91	.87	.86	.86	.88	.88
Chile ($n = 555$)	4.56	1	.033	.99	.99	.007	.080 [.019 – .161]	.95	.92	.95	.91	.80	.88	.89	.88	.88
Ecuador ($n = 477$)	.059	1	.807	1.00	1.00	.001	.000 [.000 – .076]	.91	.95	.96	.92	.84	.87	.87	.89	.89
Colombia ($n = 414$)	.454	1	.501	1.00	1.00	.002	.000 [.000 – .113]	.91	.96	.94	.93	.80	.88	.88	.89	.89
Uruguay ($n = 407$)	.346	1	.557	1.00	1.00	.003	.000 [.000 – .110]	.88	.92	.97	.84	.80	.84	.84	.84	.85
Perú ($n = 371$)	2.29	1	.130	1.00	.99	.005	.059 [.000 – .164]	.88	.95	.95	.89	.86	.86	.87	.87	.86
México ($n = 343$)	.000	1	.993	1.00	1.00	.000	.000 [.000 – .000]	.93	.92	.95	.88	.80	.87	.87	.87	.87
Cuba ($n = 342$)	1.39	1	.237	1.00	.99	.005	.034 [.000 – .153]	.91	.91	.94	.82	.77	.86	.85	.80	.81
Guatemala ($n = 344$)	1.02	1	.312	1.00	1.00	.003	.008 [.000 – .143]	.90	.95	.95	.93	.87	.88	.88	.89	.89
Argentina ($n = 316$)	.559	1	.455	1.00	1.00	.004	.000 [.000 – .134]	.95	.90	.96	.89	.81	.89	.87	.88	.87
Bolivia ($n = 264$)	.017	1	.897	1.00	1.00	.001	.000 [.000 – .074]	.87	.98	.96	.84	.84	.87	.87	.84	.84
Model 2																
Paraguay ($n = 918$)	163.35	2	.000	.99	.96	.060	.297 [.259 – .336]	.88	.88	.93	.90	–		.89	.95	
El Salvador ($n = 741$)	119.37	2	.000	.99	.96	.051	.282 [.240 – .326]	.88	.89	.91	.89	–		.90	.94	
Chile ($n = 555$)	106.38	2	.000	.99	.95	.079	.307 [.259 – .358]	.91	.90	.89	.89	–		.89	.96	
Ecuador ($n = 477$)	109.28	2	.000	.99	.96	.057	.336 [.284 – .391]	.88	.89	.93	.91	–		.90	.95	
Colombia ($n = 414$)	99.69	2	.000	.98	.95	.083	.344 [.288 – .403]	.89	.92	.90	.92	–		.89	.98	
Uruguay ($n = 407$)	64.35	2	.000	.98	.93	.067	.277 [.221 – .337]	.84	.88	.91	.83	–		.88	.92	
Perú ($n = 371$)	73.16	2	.000	.99	.96	.051	.310 [.252 – .373]	.87	.91	.90	.88	–		.89	.93	
México ($n = 343$)	79.26	2	.000	.98	.94	.070	.336 [.275 – .401]	.89	.88	.89	.87	–		.88	.94	
Cuba ($n = 342$)	73.29	2	.000	.97	.90	.073	.323 [.262 – .389]	.88	.88	.85	.78	–		.85	.91	
Guatemala ($n = 344$)	52.39	2	.000	.99	.97	.049	.271 [.211 – .337]	.88	.92	.92	.92	–		.91	.95	
Argentina ($n = 316$)	65.74	2	.000	.98	.95	.073	.318 [.255 – .386]	.92	.88	.90	.88	–		.89	.95	
Bolivia ($n = 264$)	47.31	2	.000	.99	.96	.057	.293 [.225 – .369]	.87	.93	.89	.83	–		.89	.93	

Note. ^a = Negative Error Variances/variances are negative; χ^2 = Chi square; df = degrees of freedom; SRMR: Standardized Root Mean Square Residual; TLI = Tucker-Lewis Index; CFI = Comparative Fit Index; RMSEA = Root Mean Square Error of Approximation; BIC = Sample-size adjusted Bayesian; Model 1 = Two-dimensional model; Model 2 = Unidimensional model; Φ = Relationship between dimensions; An = Anxiety dimension; De = Depression dimension



In the present study the Anxiety ($\omega = .84-.89$) and Depression ($\omega = .81-.89$) dimensions evidenced adequate internal consistency indices across countries (see Table 3). In addition, possible differences in the level of reliability between the countries were studied. The results showed no significant differences in the level of reliability in the Anxiety dimension between countries ($\chi^2(11) = 7.38$; $p = .767$). However, significant differences were evident in the level of reliability of the Depression dimension between the countries ($\chi^2(11) = 25.49$; $p = .008$).

Factor invariance by country

Table 4 shows that the factor structure of the scale has shown evidence of being strictly invariant for all countries in the sequence of invariance models proposed: metric invariance ($\Delta CFI = .000$; $\Delta RMSEA = .005$), scalar invariance ($\Delta CFI = -.001$; $\Delta RMSEA = .010$) and strict invariance ($\Delta CFI = -.002$; $\Delta RMSEA = .014$).

Table 5.

Factor invariance models of the PHQ-4 scale across countries

Invariance models	χ^2	p	TLI	CFI	SRMR	RMSEA [90%CI]	$\Delta\chi^2$	Δdf	p	ΔCFI	$\Delta RMSEA$
Configural	93.29	.001	.999	.999	.003	.038 [90%CI .024 – .052]	–	–	–	–	–
Metric	119.36	.002	.999	.999	.004	.034 [90%CI .021 – .046]	23.63	22	.367	.000	-.004
Scalar	190.09	.000	.999	.999	.005	.044 [90%CI .035 – .054]	34.72	22	.041	-.001	.010
Strict	366.09	.000	.998	.997	.014	.058 [90%CI .051 – .066]	79.03	44	.001	-.002	.014

Validity based on the relationship with other variables

Taking into account the literature review, an SEM model was proposed to evaluate the relationship between the PHQ-4 scale and well-being. It was evidenced that the structural model presents adequate fit indices ($\chi^2(24) = 960.02$; $p < .001$; $RMSEA = .084$ [CI90% .080 - .089]; $SRMR = .026$; $CFI = .99$; $TLI = .99$) and the measurement models are adequately represented by their items. Figure 1 shows that the depression dimension significantly predicts the degree of well-being ($-.66$; $p = .000$). However, the anxiety dimension fails to significantly predict the level of well-being of the participants ($-.01$; $p = .838$).




Figure 1. Predictive model of the PHQ-4 scale on the level of life satisfaction.

Discussion

According to the available literature, this is the first study to evaluate the internal structure, reliability and MI of the PHQ-4 as a screening measure of anxiety and depression symptoms in a large group of Latin American countries. Overall, the PHQ-4 demonstrated good psychometric properties in all countries; while, IM tests indicated that the factor structure and factor loadings of the PHQ-4 items were similar in all participating countries.

First, the results indicated that the PHQ-4 fits the proposed theoretical structure, since the two-factor structure presented a superior fit to the one-factor structure. This provides evidence that supports a two-dimensional model of anxious and depressive symptomatology. This finding is in agreement with previous studies reported in different cultural contexts (Ahmadi et al., 2019; Christodoulaki et al., 2022; Ghaheri et al., 2020; Kazlauskas et al., 2023; Khubchandani et al., 2016; Kim et al., 2021; Kocalevent et al., 2014; Larionow & Mudło-Głagolska, 2023; Lenz & Li, 2022; López Guerra et al., 2022; Löwe et al., 2010; Mendoza et al., 2022; Mills et al., 2015; Renovanz et al., 2019). Although the AFC indicated that the PHQ-4 has two dimensions (anxiety and depression), this finding needs further investigation, taking into account previous evidence also indicating the presence of a single factor. Regarding reliability, this was optimal as evidenced

by the values of the McDonald's ω coefficients for the anxiety and depression dimensions. This is similar to that reported by other previous psychometric studies (Ahmadi et al., 2019; Christodoulaki et al., 2022; Ghaheeri et al., 2020; Kazlauskas et al., 2023; Khubchandani et al., 2016; Kim et al., 2021; Kocalevent et al., 2014; Larionow & Mudło-Głagolska, 2023; Lenz & Li, 2022; López Guerra et al., 2022; Löwe et al., 2010; Mendoza et al., 2022; Mills et al., 2015; Renovanz et al., 2019). All this shows that the items measure both dimensions with a higher degree of accuracy.

The results of the tests of configural, metric, scalar, and strict invariance by country show that the PHQ-4 is invariant among the participating countries. Specifically, configural invariance is supported, indicating that the two-factor structure of the PHQ-4 is the same in all 12 countries. It seems that people in the 12 Latin American countries conceptualize anxiety and depression in a similar way, reflected by two factors. Likewise, the metric invariance test showed that the factor loadings were equal in all groups, which allowed us to compare the regression coefficients and covariance between different countries. In this sense, the associations between the PHQ-4 and other variables can be compared between individuals in the countries evaluated, since the change of one unit in one group is equivalent to the change in one unit in the other. Similarly, the results of the scalar invariance test show that the intersections were similar across countries, which would indicate that participants from all countries shared the same metrics and the same scale origin. Accordingly, they have the same score on the latent factors and the observed variables. This would suggest that the means of the factors can be compared between countries, as measured with the PHQ-4, and valid inferences can be made about the possible differences found (Milfont & Fischer, 2010). Finally, there was evidence of strict invariance, which indicates that the error variances are the same in all countries; therefore, the latent variables are measured with similar degrees of error between groups (van de Schoot et al., 2012). It is worth mentioning that with scalar invariance it is already possible to make comparisons between countries, however, in this study strict invariance was achieved, which ensures that the conclusions are more accurate and consistent.

The IM findings would indicate that participants in each of the countries attribute the same meaning to the latent constructs, anxiety and depression, and their scores can be compared. This is important, although the different ways in which the culture and idiosyncratic characteristics of each country can impact how people experience psychological symptoms. It has been suggested that this may be explained by the fact that the PHQ-4 items were originally selected on the basis

that they represented global symptoms of anxiety and depression that are present in different cultural contexts (Kroenke et al., 2009). Consequently, it is expected to find more similarities between the core psychological experiences, as expressed in the PHQ-4 items, than between disparate psychological experiences (Lenz & Li, 2022).

Additionally, the findings provide quantitative support for the hypothesis that the presence of depressive symptoms significantly predicts subjective well-being. However, the presence of anxiety symptoms fails to significantly predict participants' well-being. This difference has been observed before, where the association between depression and well-being holds over anxiety symptoms (Leising et al., 2013). Individuals who reported a higher frequency of depressive symptoms showed significant decreases in subjective well-being scores. It has been suggested that this finding may be explained by a shared variance in negative affect (Krieger et al., 2014). That depression, and not anxiety, significantly predicts well-being suggests that assessing depressive symptoms may have more practical utility for the assessment and development of well-being. In addition, it would allow preventive measures to be taken before the onset of depressive symptoms and help reduce healthcare costs (Fledderus et al., 2012; Grant et al., 2013; Keyes, 2007). These results provide partial support for the hypothesis of the predictive ability of depressive and anxiety symptoms on well-being.

Strengths and limitations

This study was the first to evaluate the MI of the PHQ-4 in Latin American countries and presented important strengths. First, the findings contributed to increasing the scope and application of the PHQ-4 in the Latin American context. This is important given recent evidence that the questionnaire is theoretically aligned with the measurement of symptoms of depression and anxiety (Stanhope, 2016). In addition, the study used a relatively large sample size from Latin American countries. Finally, a solid statistical analysis was performed, which took into account the ordinal nature of the PHQ-4 items.

However, the study is not free of limitations that should be considered when interpreting the results. First, the study included mostly participants from South American countries (8 of the 12 participating countries). Future studies should evaluate the MI of the PHQ-4 in even more diverse countries in the region and in other languages (such as Portuguese spoken in Brazil or Jamaican English). Second, the samples included in the study were selected by non-probability convenience sampling, which meant that the samples were not representative at the national level in each country. This limited the generalizability of the findings of the sample to the population.



Future studies should use samples obtained through probability sampling techniques to support the generalizability of the PHQ-4. Third, the samples varied in total sample size and sociodemographic characteristics (e.g., the proportion of women was higher than that of men in all countries). These differences in sociodemographic variables between country samples could further reduce the generalizability of our findings. However, previous studies established MI of the PHQ-4 between different age groups, genders, or others (Ahmadi et al., 2019; Christodoulaki et al., 2022; Ghaheri et al., 2020; Kazlauskas et al., 2023; Khubchandani et al., 2016; Kim et al., 2021; Kocalevent et al., 2014; Larionow & Mudło-Głagolska, 2023; Lenz & Li, 2022; López Guerra et al., 2022; Löwe et al., 2010; Mendoza et al., 2022; Mills et al., 2015; Renovanz et al., 2019). This would indicate that the possible confounding effects of these sociodemographic discrepancies were probably insignificant. Even so, it is recommended that future cross-national studies with the PHQ-4 employ more balanced samples with respect to sociodemographic variables in the participating countries. Fourth, an online survey was used to collect the necessary information for the study, which has the advantage of being an inexpensive and fast method, but could generate a self-selection effect of the participants, where only those who have access to the Internet could answer the questionnaire.

Fifth, suggested cut-off points for assessing IM may vary among different studies, according to different characteristics such as the type of variable or the number of participants. Although the recommendations of Chen (2007) were followed in this study, there are other more stringent criteria, such as considering a Δ CFI of 0.002 (Meade et al., 2008), or using alternative methods, such as dropping the Δ CFI to assess IM (Kang et al., 2016), which could generate different results than those reported in this study. Sixth, the use of self-report measures to assess symptoms of anxiety, depression, and subjective well-being could have generated the presence of social desirability biases in the participants. Seventh, the placement of the PHQ-4 and WHO-5 items in the online questionnaire could have been a potential limitation, because the items referring to negative emotional experiences were placed first and then the well-being items. Future studies could compare multiple versions of the online questionnaire with different item presentation orders to assess possible order effects on measurement (Rasinski et al., 2012).

Eighth, the optimal cut-off score to identify the possible presence of severe anxiety and depression was not determined. However, it should be kept in mind that the study was focused at the population level and not at the individual level, i.e., the objective was to determine the internal structure, reliability and MI of the PHQ-4 across countries. Having cut-off scores is

important to identify the number of cases with possible severe levels of anxiety and depression in a specific population, but these should be determined after having clear evidence of their internal structure, i.e., the number of dimensions assessed by the PHQ-4. In addition, the determination of optimal cut-off points requires the administration of a clinical interview, i.e., a standard measure for mental disorders. This would not be possible in population-based studies with a large sample size, as is the case in the present study. Finally, as mentioned, this study focused on the factor structure of the PHQ-4 and the MI; however, including other measures of anxiety and depression would have provided more information on the sensitivity or specificity of the PHQ-4.

Implications

The study provided evidence to consider that participants from the 12 Latin American countries perceived and interpreted the meaning of the PHQ-4 items in a similar way. Having evidence of cross-national MI of PHQ-4 scores opens new opportunities for its use in Latin America and allows for meaningful interpretations of cross-cultural comparisons of anxiety and depression symptoms between different Latin American countries and correlations with external variables. In this way, the differences in the scores of the PHQ-4 dimensions between countries are real differences and do not reflect some artificial effect associated with variations in the interpretations of the items by the participants. Furthermore, the metric invariance results give confidence to use the PHQ-4 in large epidemiological studies and to quickly determine the relationship between symptoms of anxiety, depression and other mental health indicators such as well-being within predictive models and mediation analyses. It can also be used as an early detection tool to identify groups at higher risk of experiencing high levels of anxiety and depression. This is important considering the comorbidities and long-term outcomes related to mental health symptoms in Latin America (Merchán-Hamann et al., 2012; Manrique-Millones et al., 2021; Whiteford & Baxter, 2013). The PHQ-4 can serve health professionals in different Latin American countries as a brief measure to monitor progress and assess general symptoms of depression and anxiety, as well as to evaluate whether or not an intervention has had an effect in reducing levels of anxiety and depression. Finally, the wide use of the PHQ-4 in Latin American countries may favor the development of public repositories of mental health data in this region. This is important since it is necessary to have current and regional data describing mental health disparities and needs. In addition, having this systematization of information would allow public health officials and health professionals to identify opportunities to develop initiatives that promote the wellbeing and development of Latin Americans.



Conclusion

As mentioned above, this study represented a first attempt to evaluate the MI of the PHQ-4 cross-culturally in the Spanish language and in the Latin American context. Despite the limitations, it is concluded that the PHQ-4 provides a largely comparable measure of anxiety and depressive symptoms in a large sample of Latin American countries. This provides preliminary and promising evidence for the use of the PHQ-4 in a broader context of primary care practice and research activities.

References

- Ahmadi, S. M., Arani, A. M., Bakhtiari, M., & Emamy, M. H. D. (2019). Psychometric properties of Persian version of patient health questionnaires-4 (PHQ-4) in coronary heart disease patients. *Iranian Journal of Psychiatry and Behavioral Sciences*, 13(4), e85820. <https://doi.org/10.5812/ijpbs.85820>
- Ali, A. M., Hori, H., Kim, Y., & Kunugi, H. (2022). The Depression Anxiety Stress Scale 8-items expresses robust psychometric properties as an ideal shorter version of the Depression Anxiety Stress Scale 21 among healthy respondents from three continents. *Frontiers in Psychology*, 13, 799769. <https://doi.org/10.3389/fpsyg.2022.799769>
- Baumgartner, H., & Steenkamp, J. B. E. (1998). Multi-group latent variable models for varying numbers of items and factors with cross-national and longitudinal applications. *Marketing Letters*, 9, 21-35. <https://doi.org/10.1023/A:1007911903032>
- Beck, A. T., Ward, C. H., Mendelson, M., Mock, J., & Erbaugh, J. (1961). An inventory for measuring depression. *Archives of General Psychiatry*, 4(6), 561-571. <https://doi.org/10.1001/archpsyc.1961.01710120031004>
- Boer, D., Hanke, K., & He, J. (2018). On detecting systematic measurement error in cross-cultural research: A review and critical reflection on equivalence and invariance tests. *Journal of Cross-Cultural Psychology*, 49(5), 713-734. <https://doi.org/10.1177/0022022117749042>
- Bowden, A., & Fox-Rushby, J. A. (2003). A systematic and critical review of the process of translation and adaptation of generic health-related quality of life measures in Africa, Asia, Eastern Europe, the Middle East, South America. *Social Science & Medicine*, 57(7), 1289-1306. [https://doi.org/10.1016/S0277-9536\(02\)00503-8](https://doi.org/10.1016/S0277-9536(02)00503-8)
- Byrne, B. M. (2016). Multigroup comparisons: Testing for measurement, structural, and latent mean equivalence. In F. T. L. Leong, D. Bartram, F. M. Cheung, K. F. Geisinger, & D. Iliescu (Eds.), *The ITC international handbook of testing and assessment* (pp. 377–394). Oxford University Press. <https://doi.org/10.1093/med:psych/9780199356942.003.0026>
- Caldas de Almeida, J. M., & Horvitz-Lennon, M. (2010). Mental health care reforms in Latin America: an overview of mental health care reforms in Latin America and the Caribbean. *Psychiatric Services*, 61(3), 218-221. <https://doi.org/10.1176/ps.2010.61.3.218>

- Caycho-Rodríguez, T., Vilca, L. W., Valencia, P. D., Carbajal-León, C., Reyes-Bossio, M., White, M., ... & Flores-Mendoza, C. (2023). Is the meaning of subjective well-being similar in Latin American countries? A cross-cultural measurement invariance study of the WHO-5 well-being index during the COVID-19 pandemic. *BMC Psychology*, 11(1), 1-14. <https://doi.org/10.1186/s40359-023-01149-8>
- Cenat, J. M., Farahi, S. M. M. M., Dalexis, R. D., Darius, W. P., Bekarkhanechi, F. M., Poisson, H., ... & Labelle, P. R. (2022). The global evolution of mental health problems during the COVID-19 pandemic: A systematic review and meta-analysis of longitudinal studies. *Journal of Affective Disorders*, 315, 70-95. <https://doi.org/10.1016/j.jad.2022.07.011>
- Chen, W. H., & Thissen, D. (1997). Local dependence indexes for item pairs using item response theory. *Journal of Educational Behavioral Statistics*, 22, 265–289. <https://doi.org/10.2307/1165285>
- Christodoulaki, A., Baralou, V., Konstantakopoulos, G., & Touloumi, G. (2022). Validation of the Patient Health Questionnaire-4 (PHQ-4) to screen for depression and anxiety in the Greek general population. *Journal of Psychosomatic Research*, 160, 110970. <https://doi.org/10.1016/j.jpsychores.2022.110970>
- Clark, D. M. (2011). Implementing NICE guidelines for the psychological treatment of depression and anxiety disorders: the IAPT experience. *International Review of Psychiatry*, 23(4), 318-327. <https://doi.org/10.3109/09540261.2011.606803>
- Cronbach, L. J. (1951). Coefficient alpha and the internal structure of tests. *Psychometrika*, 16(3), 297-334. <https://doi.org/10.1007/BF02310555>
- Davidov, E., Meuleman, B., Cieciuch, J., Schmidt, P., & Billiet, J. (2014). Measurement equivalence in cross-national research. *Annual Review of Sociology*, 40, 55-75. <https://doi.org/10.1146/annurev-soc-071913-043137>
- Diedenhofen, B., & Musch, J. (2016). cocron: A Web Interface and R Package for the Statistical Comparison of Cronbach's Alpha Coefficients. *International Journal of Internet Science*, 11(1), 51-60
- Errazuriz, A., & Crisostomo, N. (2021). Prevalence of depression in Latin America and the Caribbean: protocol for a systematic review and meta-analysis. *JBI Evidence Synthesis*, 19(1), 201-207. <https://doi.org/10.11124/JBISRIR-D-19-00364>
- Eysenbach, G. (2004). Improving the quality of Web surveys: The Checklist for Reporting Results of Internet E-Surveys (CHERRIES). *Journal of Medical Internet Research*, 6(3), e34. <https://doi.org/10.2196/jmir.6.3.e34>
- Feldt, L. S., Woodruff, D. J., & Salih, F. A. (1987). Statistical inference for coefficient alpha. *Applied Psychological Measurement*, 11(1), 93-103. <https://doi.org/10.1177/014662168701100107>
- Fernández, R. S., Crivelli, L., Guimet, N. M., Allegri, R. F., Picco, S., & Pedreira, M. E. (2022). Psychological distress and mental health trajectories during the COVID-19 pandemic in



- Argentina: a longitudinal study. *Scientific Reports*, 12(1), 5632. <https://doi.org/10.1038/s41598-022-09663-2>
- Fledderus, M., Bohlmeijer, E. T., Pieterse, M. E., & Schreurs, K. M. G. (2012). Acceptance and commitment therapy as guided self-help for psychological distress and positive mental health: a randomized controlled trial. *Psychological Medicine*, 42(3), 485-495. <https://doi.org/10.1017/S0033291711001206>
- Fong, T. C., Ho, R. T., & Yip, P. S. (2023). Psychometric properties of the Patient Health Questionnaire-4 among Hong Kong young adults in 2021: Associations with meaning in life and suicidal ideation. *Frontiers in Psychiatry*, 14, 1138755. <https://doi.org/10.3389/fpsy.2023.1138755>
- Ghaheri, A., Omani-Samani, R., Sepidarkish, M., Hosseini, M., & Maroufizadeh, S. (2020). The four-item patient health questionnaire for anxiety and depression: A validation study in infertile patients. *International Journal of Fertility & Sterility*, 14(3), 234-239. <https://doi.org/10.22074/ijfs.2020.44412>
- Gilbody, S. M., House, A. O., & Sheldon, T. A. (2002). Psychiatrists in the UK do not use outcomes measures: National survey. *The British Journal of Psychiatry*, 180(2), 101-103. <https://doi.org/10.1192/bjp.180.2.101>
- Grant, F., Guille, C., & Sen, S. (2013). Well-being and the risk of depression under stress. *PLoS one*, 8(7), e67395. <https://doi.org/10.1371/journal.pone.0067395>
- Hinz, A., Finck, C., Gómez, Y., Daig, I., Glaesmer, H., & Singer, S. (2014). Anxiety and depression in the general population in Colombia: reference values of the Hospital Anxiety and Depression Scale (HADS). *Social Psychiatry and Psychiatric Epidemiology*, 49, 41-49. <https://doi.org/10.1007/s00127-013-0714-y>
- Horn, J. L., & McArdle, J. J. (1992). A practical and theoretical guide to measurement invariance in aging research. *Experimental Aging Research*, 18(3), 117-144. <https://doi.org/10.1080/03610739208253916>
- Jorgensen, T. D., Pornprasertmanit, S., Schoemann, A. M., & Rosseel, Y. (2018). *semTools: Useful tools for structural equation modeling. R package version 0.5-1*. <https://CRAN.R-project.org/package=semTools>
- Kang, Y., McNeish, D. M., & Hancock, G. R. (2016). The role of measurement quality on practical guidelines for assessing measurement and structural invariance. *Educational and Psychological Measurement*, 76(4), 533-561. <https://doi.org/10.1177/0013164415603764>
- Katon, W., & Roy-Byrne, P. (2007). Anxiety disorders: efficient screening is the first step in improving outcomes. *Annals of Internal Medicine*, 146(5), 390-392. <https://doi.org/10.7326/0003-4819-146-5-200703060-00011>
- Kazlauskas, E., Gelezelyte, O., Kvedaraitė, M., Ajdukovic, D., Johannesson, K. B., Böttche, M., ... & Lotzin, A. (2023). Psychometric properties of the Patient Health Questionnaire-4 (PHQ-4) in 9230 adults across seven European countries: Findings from the ESTSS

- ADJUST study. *Journal of Affective Disorders*, 335, 18-23. <https://doi.org/10.1016/j.jad.2023.05.007>
- Keyes, C. L. M. (2007). Promoting and protecting mental health as flourishing: A complementary strategy for improving national mental health. *American Psychologist*, 62(2), 95–108. <https://doi.org/10.1037/0003-066X.62.2.95>
- Khubchandani, J., Brey, R., Kotecki, J., Kleinfelder, J., & Anderson, J. (2016). The psychometric properties of PHQ-4 depression and anxiety screening scale among college students. *Archives of Psychiatric Nursing*, 30(4), 457-462. <https://doi.org/10.1016/j.apnu.2016.01.014>
- Kim, H. W., Shin, C., Lee, S. H., & Han, C. (2021). Standardization of the Korean version of the Patient Health Questionnaire-4 (PHQ-4). *Clinical Psychopharmacology and Neuroscience*, 19(1), 104–111. <https://doi.org/10.1007/s12035-021-01104-1>
- Kinderman, P., Tai, S., Pontin, E., Schwannauer, M., Jarman, I., & Lisboa, P. (2015). Causal and mediating factors for anxiety, depression and well-being. *The British Journal of Psychiatry*, 206(6), 456-460. <https://doi.org/10.1192/bjp.bp.114.147553>
- Kliem, S., Mößle, T., Klatt, T., Fleischer, S., Kudlacek, D., Kröger, C., ... & Wiltink, J. (2016). Psychometrische Prüfung einer Hocharabischen Übersetzung des PHQ-4 anhand einer repräsentativen Befragung syrischer Geflüchteter [Psychometric Testing of a Standard Arabic Translation of the PHQ-4 Using a Representative Survey of Syrian Refugees]. *PPmP-Psychotherapie· Psychosomatik· Medizinische Psychologie*, 66(09/10), 385-392. <https://doi.org/10.1055/s-0042-114775>
- Kline, R. B. (2015). *Principles and Practice of Structural Equation Modeling* (4th ed.). The Guilford Press.
- Kocalevent, R. D., Finck, C., Jimenez-Leal, W., Sautier, L., & Hinz, A. (2014). Standardization of the Colombian version of the PHQ-4 in the general population. *BMC Psychiatry*, 14(1), 1-8. <https://doi.org/10.1186/1471-244X-14-205>
- Kohn, R., Ali, A. A., Puac-Polanco, V., Figueroa, C., López-Soto, V., Morgan, K., ... & Vicente, B. (2018). Mental health in the Americas: an overview of the treatment gap. *Revista Panamericana de Salud Pública*, 42, e165. <https://doi.org/10.26633/RPSP.2018.165>
- Krieger, T., Zimmermann, J., Huffziger, S., Ubl, B., Diener, C., Kuehner, C., & Holtforth, M. G. (2014). Measuring depression with a well-being index: further evidence for the validity of the WHO Well-Being Index (WHO-5) as a measure of the severity of depression. *Journal of Affective Disorders*, 156, 240-244. <https://doi.org/10.1016/j.jad.2013.12.015>
- Kroenke, K., Spitzer, R. L., Williams, J. B., & Löwe, B. (2009). An ultra-brief screening scale for anxiety and depression: the PHQ-4. *Psychosomatics*, 50(6), 613-621. [https://doi.org/10.1016/S0033-3182\(09\)70864-3](https://doi.org/10.1016/S0033-3182(09)70864-3)
- Kroenke, K., Spitzer, R. L., Williams, J. B., Monahan, P. O., & Löwe, B. (2007). Anxiety disorders in primary care: prevalence, impairment, comorbidity, and detection. *Annals of Internal Medicine*, 146(5), 317-325. <https://doi.org/10.7326/0003-4819-146-5-200703060-00004>



- Lahlouh, A., & Mustafa, M. (2020). Sleep quality and health related problems of shift work among resident physicians: a cross-sectional study. *Sleep Medicine*, 66, 201-206. <https://doi.org/10.1016/j.sleep.2019.11.1258>
- Larionow, P., & Mudło-Głagolska, K. (2023). The Patient Health Questionnaire–4: Factor Structure, Measurement Invariance, Latent Profile Analysis of Anxiety and Depressive Symptoms and Screening Results in Polish Adults. *Advances in Cognitive Psychology*, 19(2), 123-137. <https://doi.org/10.5709/acp-0391-2>
- Leising, D., Borkenau, P., Zimmermann, J., Roski, C., Leonhardt, A., & Schütz, A. (2013). Positive self-regard and claim to leadership: Two fundamental forms of self-evaluation. *European Journal of Personality*, 27(6), 565-579. <https://doi.org/10.1002/per.1924>
- Lenz, A. S., & Li, C. (2022). Evidence for measurement invariance and psychometric reliability for scores on the PHQ-4 from a rural and predominately hispanic community. *Measurement and Evaluation in Counseling and Development*, 55(1), 17-29. <https://doi.org/10.1080/07481756.2021.1906157>
- Li, C. H. (2016). Confirmatory factor analysis with ordinal data: Comparing robust maximum likelihood and diagonally weighted least squares. *Behavior Research Methods*, 48, 936-949. <https://doi.org/10.3758/s13428-015-0619-7>
- López Guerra, V., Aguirre Mejia, Á. J., & Guerrero Alcedo, J. M. (2022). Propiedades psicométricas y estructura factorial del cuestionario de salud del paciente PHQ-4 en estudiantes universitarios ecuatorianos [Psychometric properties and factor structure of the PHQ-4 patient health questionnaire in Ecuadorian university students] . *Revista Cubana de Enfermería*, 38(3), e4885. <http://scielo.sld.cu/pdf/enf/v38n3/1561-2961-enf-38-03-e4885.pdf>
- Lopez, A. O., Martinez, M. N., Garcia, J. M., Kunik, M. E., & Medina, L. D. (2021). Self-report depression screening measures for older Hispanic/Latin American adults: A PRISMA systematic review. *Journal of Affective Disorders*, 294, 1-9. <https://doi.org/10.1016/j.jad.2021.06.049>
- López-Rodríguez, J. A. (2019). Declaración de la iniciativa CHERRIES: adaptación al castellano de directrices para la comunicación de resultados de cuestionarios y encuestas online [CHERRIES Initiative Statement: Adaptation into Spanish of guidelines for communicating the results of online questionnaires and surveys]. *Atención Primaria*, 51(9), 586-589. <https://doi.org/10.1016/j.aprim.2019.03.005>
- Löwe, B., Kroenke, K., & Gräfe, K. (2005). Detecting and monitoring depression with a two-item questionnaire (PHQ-2). *Journal of Psychosomatic Research*, 58(2), 163-171. <https://doi.org/10.1016/j.jpsychores.2004.09.006>
- Löwe, B., Wahl, I., Rose, M., Spitzer, C., Glaesmer, H., Wingenfeld, K., ... & Brähler, E. (2010). A 4-item measure of depression and anxiety: validation and standardization of the Patient Health Questionnaire-4 (PHQ-4) in the general population. *Journal of Affective Disorders*, 122(1-2), 86-95. <https://doi.org/10.1016/j.jad.2009.06.019>

- Malone, C., & Wachholtz, A. (2018). The relationship of anxiety and depression to subjective well-being in a mainland Chinese sample. *Journal of Religion and Health, 57*, 266-278. <https://doi.org/10.1007/s10943-017-0447-4>
- Manrique-Millones, D., Wium, N., Pineda-Marín, C., Fernández-Arata, M., Alfonso-Murcia, D., López-Martínez, J. L., & Millones-Rivalles, R. (2021). Association between substance use behaviors, developmental assets and mental health: a glance at Latin American young college students. *Frontiers in Psychology, 12*, 639578. <https://doi.org/10.3389/fpsyg.2021.639578>
- Materu, J., Kuringe, E., Nyato, D., Galishi, A., Mwanamsangu, A., Katebalila, M., ... & Wambura, M. (2020). The psychometric properties of PHQ-4 anxiety and depression screening scale among out of school adolescent girls and young women in Tanzania: a cross-sectional study. *BMC Psychiatry, 20*(1), 1-8. <https://doi.org/10.1186/s12888-020-02735-5>
- Mautong, H., Gallardo-Rumbea, J. A., Alvarado-Villa, G. E., Fernández-Cadena, J. C., Andrade-Molina, D., Orellana-Román, C. E., & Cherrez-Ojeda, I. (2021). Assessment of depression, anxiety and stress levels in the Ecuadorian general population during social isolation due to the COVID-19 outbreak: a cross-sectional study. *BMC Psychiatry, 21*(1), 1-15. <https://doi.org/10.1186/s12888-021-03214-1>
- McDonald, R. P. (1999). *Test Theory: A Unified Treatment*. Taylor & Francis.
- Meade, A. W., Johnson, E. C., & Braddy, P. W. (2008). Power and sensitivity of alternative fit indices in tests of measurement invariance. *Journal of Applied Psychology, 93*(3), 568–592. <https://doi.org/10.1037/0021-9010.93.3.568>
- Mendoza, N. B., Frondoza, C. E., Dizon, J. I. W. T., & Buenconsejo, J. U. (2022). The factor structure and measurement invariance of the PHQ-4 and the prevalence of depression and anxiety in a Southeast Asian context amid the COVID-19 pandemic. *Current Psychology, 1-10*. <https://doi.org/10.1007/s12144-022-02833-5>
- Merchán-Hamann, E., Leal, E. M., Basso Musso, L., García Estrada, M., Reid, P., Kulakova, O. V., ... & Domenech, D. (2012). Comorbilidad entre abuso/dependencia de drogas y el distrés psicológico en siete países de Latinoamérica y uno del Caribe [Comorbidity between drug abuse/dependence and psychological distress in seven Latin American countries and one Caribbean country]. *Texto & Contexto-Enfermagem, 21*, 87-95. <https://doi.org/10.1590/S0104-07072012000500012>
- Milfont, T. L., & Fischer, R. (2010). Testing measurement invariance across groups: Applications in cross-cultural research. *International Journal of Psychological Research, 3*(1), 111-130.
- Mills, S. D., Fox, R. S., Pan, T. M., Malcarne, V. L., Roesch, S. C., & Sadler, G. R. (2015). Psychometric evaluation of the patient health questionnaire–4 in Hispanic Americans. *Hispanic Journal of Behavioral Sciences, 37*(4), 560-571. <https://doi.org/10.1177/0739986315608126>
- Pan American Health Organization (2023). *Salud Mental*. <https://www.paho.org/es/temas/salud-mental>

- Parkitny, L., & McAuley, J. (2010). The depression anxiety stress scale (DASS). *Journal of Physiotherapy*, 56(3), 204.
- R Core Team. (2019). *A language and environment for statistical computing (R version 3.6.1)*. R Foundation for Statistical Computing. <http://www.r-project.org/>
- Rasinski, K. A., Lee, L., & Krishnamurty, P. (2012). Question order effects. In H. Cooper, P. M. Camic, D. L. Long, A. T. Panter, D. Rindskopf, & K. J. Sher (Eds.), *APA handbook of research methods in psychology, Vol. 1. Foundations, planning, measures, and psychometrics* (pp. 229–248). American Psychological Association. <https://doi.org/10.1037/13619-014>
- Renovanz, M., Soebianto, S., Tsakmaklis, H., Keric, N., Nadji-Ohl, M., Beutel, M., ... & Hickmann, A. K. (2019). Evaluation of the psychological burden during the early disease trajectory in patients with intracranial tumors by the ultra-brief Patient Health Questionnaire for Depression and Anxiety (PHQ-4). *Supportive Care in Cancer*, 27, 4469-4477. <https://doi.org/10.1007/s00520-019-04718-z>
- Rosseel, Y. (2012). lavaan : An R Package for Structural Equation Modeling. *Journal of Statistical Software*, 48(2), 1–36. <https://doi.org/10.18637/jss.v048.i02>
- RStudio Team. (2018). *RStudio: Integrated Development Environment for R*. RStudio, Inc. <http://www.rstudio.com/>
- Ruiz, F. J., Martín, M. B. G., Falcón, J. C. S., & González, P. O. (2017). The hierarchical factor structure of the Spanish version of Depression Anxiety and Stress Scale-21. *International Journal of Psychology and Psychological Therapy*, 17(1), 97-105.
- Schumacker, R. E., & Lomax, R. G. (2015). *A Beginner's Guide to Structural Equation Modeling: Fourth Edition* (4th ed.). Routledge.
- Soper, D.S. (2023). *A-priori sample size calculator for structural equation models* [Software]. <http://www.danielsoper.com/statcalc>
- Stanhope, J. (2016). Patient health questionnaire-4. *Occupational Medicine*, 66(9), 760-761. <https://doi.org/10.1093/occmed/kqw165>
- Topp, C. W., Østergaard, S. D., Søndergaard, S., & Bech, P. (2015). The WHO-5 Well-Being Index: a systematic review of the literature. *Psychotherapy and Psychosomatics*, 84(3), 167-176. <https://doi.org/10.1159/000376585>
- Van de Schoot, R., Lugtig, P., & Hox, J. (2012). A checklist for testing measurement invariance. *European Journal of Developmental Psychology*, 9(4), 486-492. <https://doi.org/10.1080/17405629.2012.686740>
- van de Vijver, F., & Tanzer, N. K. (2004). Bias and equivalence in cross-cultural assessment: An overview. *European Review of Applied Psychology*, 54(2), 119-135. <https://doi.org/10.1016/j.erap.2003.12.004>
- Viladrich, C., Angulo-Brunet, A., & Doval, E. (2017). A journey around alpha and omega to estimate internal consistency reliability. *Anales de Psicología*, 33(3), 755–782.

<https://doi.org/10.6018/analesps.33.3.268401>

Whiteford, H. A., & Baxter, A. J. (2013). The Global Burden of Disease 2010 Study: what does it tell us about mental disorders in Latin America? *Brazilian Journal of Psychiatry*, 35, 111-112. <https://doi.org/10.1590/1516-4446-2012-3502>

Wicke, F. S., Krakau, L., Löwe, B., Beutel, M. E., & Brähler, E. (2022). Update of the standardization of the Patient Health Questionnaire-4 (PHQ-4) in the general population. *Journal of Affective Disorders*, 312, 310-314. <https://doi.org/10.1016/j.jad.2022.06.054>

World Health Organization (1998). *Well-Being measures in primary health care: The DepCare Project. Health for All. Target 12*. Geneva, WHO.

World Health Organization (2022). *La OMS subraya la urgencia de transformar la salud mental y los cuidados conexos* [WHO stresses urgency to transform mental health and related care]. <https://www.who.int/es/news/item/17-06-2022-who-highlights-urgent-need-to-transform-mental-health-and-mental-health-care>

Zhang, S. X., Batra, K., Xu, W., Liu, T., Dong, R. K., Yin, A., ... & Chen, J. (2022). Mental disorder symptoms during the COVID-19 pandemic in Latin America—a systematic review and meta-analysis. *Epidemiology and Psychiatric Sciences*, 31, e23. <https://doi.org/10.1017/S2045796021000767>

Zigmond, A. S., & Snaith, R. P. (1983). The hospital anxiety and depression scale. *Acta Psychiatrica Scandinavica*, 67(6), 361-370. <https://doi.org/10.1111/j.1600-0447.1983.tb09716.x>

About the Authors

Tomás Caycho-Rodríguez, Ph.D. in Psychology. Senior Researcher at the Universidad Científica del Sur, Research Renacyt Distinguished category. His research interests are psychometrics, psychogerontology, cross-cultural research.

Lindsey W. Vilca, Master in Psychology. Professor and researcher at the Norbert Wiener University in Lima, Peru. His research interests are psychometrics and clinical and health Psychology.

José Ventura-León, Ph.D. in Psychology. Senior Researcher at the Universidad Privada del Norte, Renacyt Distinguished Researcher. His research interests are psychometrics, network analysis, statistics, clinical psychology.

Pablo D. Valencia, PhD Student, School of Higher Studies (F.E.S.) Iztacala, National Autonomous University of Mexico.



Carlos Carbajal-León, Ph.D. in psychology and Master in Educational Psychology. Research professor at the Peruvian University of Applied Sciences.

Mario Reyes-Bossio, Master in Psychology, University of San Martín de Porres. Researcher at the Peruvian University of Applied Sciences.

Mariel Delgado-Campusano, Master in Psychology, Researcher at the Peruvian University of Applied Sciences.

Daniel E. Yupanqui-Lorenzo, Researcher at the University of Sciences and Humanities.

Claudio Rojas-Jara, Master in drug addiction. Master in prevention and treatment of addictive behaviors. Academic of the Department of Psychology, Faculty of Health Sciences, Universidad Católica del Maule, Chile.

Miguel Gallegos, Ph.D. in Psychology from the National University of Rosario, Argentina, and Doctor of Education from the Federal University of Minas Gerais, Brazil. Post-doctorate from the National Autonomous University of Mexico, and the Pontificia Universidade Católica de Minas Gerais, Brazil.

Mauricio Cervigni, Ph.D. in Psychology from the National University of Rosario (UNR). Researcher at the National Council for Scientific and Technical Research (CONICET - Argentina). Professor-Researcher at the National University of Rosario (UNR).

Pablo Martino, Ph.D. in Psychology (National University of San Luis-Arg). Adjunct professor at the Faculty of Psychology, National University of Rosario. National Council for Scientific and Technical Research postdoctoral fellow.

Roberto Polanco-Carrasco, Bachelor of Psychology, Scientific Editor of Cuadernos de Neuropsicología – Panamerican Journal of Neuropsychology. Director of the Chilean Association of Scientific Journals of Psychology.

Diego Alejandro Palacios, Clinical Psychologist, Master in Clinical Psychology and Mental Health. Psychotherapist at the Human Development Center Clinics -CDH- and teacher at the University of San Carlos de Guatemala.

Rodrigo Moreta-Herrera, Clinical Psychologist and University Master's Degree in Psychology. Doctor (c) in Psychology from the University of Girona. Professor and researcher at the Pontifical Catholic University of Ecuador.

Antonio Samaniego-Pinho, Psychologist with a specialty in Occupational Psychology. Master in Clinical Psychology. Teacher in the career of Psychology at the Faculty of Philosophy (National University of Asuncion).

Marlon Elías Lobos Rivera, Psychologist. Master in University Education. Research profesor at the School of Psychology, Faculty of Social Sciences, Technological University of El Salvador.

Andrés Buschiazco Figares, Degree in Psychology (University of the Republic, Uruguay). Psychotherapist. Academic Director of the Center for Adlerian Studies, Uruguay.

Diana Ximena Puerta-Cortés, Ph.D. in Psychology (Ramon LLull University of Barcelona, Spain). Master's degree in psychology with an emphasis on addictions and violence (Universidad Católica de Colombia). She is a researcher for the GESS group and director of the NeuroTech research hotbed. Professor at the University of Ibagué in Colombia.

Ibraín Enrique Corrales-Reyes, Dentist. Maxillofacial surgeon. Professor and researcher at the Carlos Manuel de Céspedes General University Hospital. Granma University of Medical Sciences. Cuba. Editor of the Cuban Journal of Stomatology and the Cuban Journal of Biomedical Research.

Raymundo Calderón, Master of Science in Education. National Director of Psychology, Universidad Del Valle de México.

Olimpia Petzold, Ph.D. in Psychology, Lone Star College, Conroe, The United States of America.

Andrés Camargo, Ph.D. in Psychology, School of Health and Sport Sciences, Fundación Universitaria del Area Andina, Bogotá, Colombia.

Julio Torales, Head of the Neuroscience Department. School of Medical Sciences, Universidad Nacional de Asunción.

Daniela Ferrufino-Borja, Graduate in Psychology, Psychological Research and Counseling Center, Faculty of Humanities, Communication and Arts, Private University of Santa Cruz de la Sierra, Santa Cruz, Bolivia.

Agueda Muñoz-del-Carpio-Toia, Doctor of Medicine and Master of Public Health. Full-time Research Professor in the Vice-Rector's Office for Research. Professor at the Graduate School of the Catholic University of Santa María (UCSM).



Jorge Palacios, Ph.D. in Psychology, Professor of Psychology, Faculty of Health Sciences, Universidad del Valle de Mexico, Mexico City, Mexico.

Marion K. Schulmeyer, Graduate in Philosophy and Letters, Psychology Section from the Universidad Pontificia Comillas (Spain). Dean of the Faculty of Humanities and Communication of the Private University of Santa Cruz de la Sierra, Santa Cruz, Bolivia.

Cirilo H. Garcia-Cadena, PhD in Social Policy from the University of Texas. Professor and researcher at the Faculty of Psychology of the Autonomous University of Nuevo León. Organizational consultant at the Faculty of Psychology of the Autonomous University of Nuevo León.

María José Baeza-Rivera, Psychologist from the University of La Frontera, Chile. PhD in Psychology, University of La Frontera.

Camila Salazar-Fernández, PhD in Psychology. Professor at the Department of Psychology at the University of La Frontera.

Nicol A. Barria-Asenjo, Researcher in Social Sciences and Humanities. Author of essays, books and academic articles published in specialized/academic journals and publishers. Editor of special issues of scientific and academic journals in Latin America and Europe.

Corresponding Author`s Address ^[TOP]

Facultad de Psicología, Universidad Científica del Sur

Campus Villa II, Ctra. Panamericana S 19, Villa EL Salvador, Lima, Peru

Email: tcaycho@cientifica.edu.pe